PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Docket No: Q386121

RECEIVED

Munchisa FUJITA, et al.

OCT 2 4 2003

Appln. No.: 08/915,683

Group Art Unit: 1752

TC 1700

Confirmation No.: 1847

Examiner: WALKE, Amanda C.

Filed: August 21, 1997

For:

DIRECT-POSITIVE PHOTOGRAPHIC SILVER HALIDE EMULSION AND COLOR PHOTOGRAPHIC LIGHT-SENSITIVE MATERIAL COMPRISING SAME

FOURTH SUPPLEMENTAL DECLARATION UNDER 37 C.F.R. § 1.132

Commissioner for Patents Washington, D.C. 20231

Sir:

I, Takefurni HARA, do hereby declare and state as follows:

I am the same declarant who performed the experimentation described in the Third Supplemental Declaration Under 37 C.F.R. § 1.132 dated April 21, 2003, and filed on April 23, 2003. My personal history remains as stated in my prior Declaration.

The data and explanation presented in this Fourth Supplemental Declaration demonstrate that the effect of decreasing the reversal negative sensitivity due to the emulsion of the present invention is unexpectedly large.

As set forth in the Table below, I compare the density of negative image corresponding to a reversal negative relative sensitivity of 88 or less with the density of negative image corresponding to a reversal negative relative sensitivity of larger than 90.

FOURTH SUPPLEMENTAL DECLARATION UNDER 37 C.F.R. § 1.132 U.S. Appln. No. 08/915,683

When a sample having a negative image relative sensitivity of 100 is exposed to light such that the density of the negative image is 1.50, the densities of the negative images of samples having various relative sensitivities are shown below. Such difference in density is significant and can be visually observed by naked eyes of human.

Relative sensitivity of negative image of sample	Density of negative image		
100	1.50		
95	1.30 1.12 1.05 0.95		
90			
88			
85			
80	0.78		

In the Third Supplemental Declaration, Sample Nos. 25 to 28 were prepared in the same procedure as Sample Nos. 13-16, except that Compound B which contains no sulfur atom, was used as a solvent, instead of Compound A (a sulfur-containing solvent). Sample Nos. 25 to 28 were then compared with Sample Nos. 13-16. The results are summarized in the table below:

FOURTH SUPPLEMENTAL DECLARATION UNDER 37 C.F.R. § 1.132 U.S. Appln. No. 08/915.683

Sample No.	Y M C	Y M C	Middle sensitivity	Negative sensitivity	Coefficien of variation
25 (inv)	2.17 2.31 2.47		Y M C	Y M C	in grain thickness
13 (inv)		1.20 0.24	110 126 147	83 83 83	16%
26 (inv)	2.18 2.32 2.48	0.15 0.16 0.23	111 128 149	82 82 81	15%
14 (inv)	2.13 2.27 2.43	0.17 0.18 0.26	108 124 146	86 88 88	29%
27 (comp)	2.14 2.29 2.45	0.17 0.18 0.26	109 125 147	85 88 87	28%
	2.10 2.24 2.40	0.19 0.21 0.29	105 121 142	91 93 94	35%
15 (comp)	2.12 2.26 2.42	0.19 0.20 0.28	105 120 141	92-94-95	-
28 (comp)	2.04 2.19 2.33	0.23 0.24 0.33	10:	100 103 105	33%
16 (comp)	2.06 2.21 2.35	0.23 0.24 0.32	102 119 138		45%
		0.55	102 119 138	99 101 103	43%

As is clear from the above results, substantially the same results were obtained in Sample Nos. 25-28 wherein non-sulfur atom-containing compound B was used as a solvent and Sample Nos. 13-16 wherein sulfur atom-containing compound A was used as a solvent. That is, compound A, as a solvent, does not have any significant effect on the photographic properties of an emulsion.

Further, Sample Nos. 25 and 26 of the present invention were higher in D_{max} and middle sensitivity, but remarkably lower in D_{min} and Negative sensitivity than Comparative Sample Nos. 27 and 28. Specifically, Sample Nos. 25 and 26 have a negative of sensitivity of between 81 to 88, whereas Sample Nos. 27 and 28 have a negative sensitivity of between 91 to 105. As explained above, this difference in the negative sensitivity is significant, in that the difference in the densities of negative images after the samples are exposed to light is significant.

FOURTH SUPPLEMENTAL DECLARATION UNDER 37 C.F.R. § 1.132 U.S. Appin. No. 08/915,683

The above experimentation concretely demonstrates that the unexpected results of the present invention are obtained independent of the presence of sulfur atom in the silver halide solvent, and further, that control of coefficient of variation of grain thickness is critical to achieving the effects of the present invention.

In view of the above, I conclude that the effect of decreasing the reversal negative sensitivity due to the emulsion of the present-invention, having a coefficient of variation in grain thickness of not more than 30%, is significant and unexpected.

I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: October 7, 2003

Jakefuni Hara Takefumi HARA